

- TINBERGEN, L. 1960. The natural control of insects in pine-woods. I: Factors influencing the intensity of predation by songbirds. *Arch. Neerl. Zool.* 13:266–336.
- TURNER, J. R. G. 1984. Mimicry: the palatability spectrum and its consequences, pp. 141–161. *In* Vane-Wright, R. I. & P. R. Ackery (eds.), *The biology of butterflies*, 2nd ed. Princeton Univ. Press, Princeton, New Jersey.
- TYLER, H. A., K. S. BROWN JR. & K. H. WILSON. 1994. *Swallowtail butterflies of the Americas*. Scientific Publishers, Gainesville, Florida.
- VAUGHAN, F. A. 1983. Startle responses of blue jays to visual stimuli presented during feeding. *Anim. Behav.* 31:385–396.
- WENZEL, J. W. 1992. Behavioral homology and phylogeny. *Ann. Rev. Ecol. Syst.* 23:361–381.
- WIKLUND, C. & T. JÄRVI. 1982. Survival of distasteful insects after being attacked by naive birds: a reappraisal of aposematic coloration evolving through individual selection. *Evolution* 36:998–1002.
- WIKLUND, C. & B. SILLÉN-TULLBERG. 1985. Why distasteful butterflies have aposematic larvae and adults, but cryptic pupae: evidence from predation experiments on the monarch and the European swallowtail. *Evolution* 39:1155–1158.
- WILLIAMS, G. C. 1966. *Adaptation and natural selection*. Princeton Univ. Press, Princeton, New Jersey.

ANDREW V. Z. BROWER, *Department of Entomology, Oregon State University, Corvallis, Oregon 97331, USA*, AND KAREN R. SIME, *Department of Entomology & Section of Ecology and Systematics, Cornell University, Ithaca, New York 14853, USA*.

Received for publication 6 December 1996; revised and accepted 28 August 1997.

Journal of the Lepidopterists' Society
52(2), 1998, 212–214

FIELD OBSERVATIONS ON MATING BEHAVIOR AND PREDATION OF *HEMILEUCA ELECTRA* (SATURNIIDAE)

Additional key words: visual cues, predation, silk moth, mate location, pheromones.

Hemileuca electra (Wright), the Electra Buckmoth, is a widespread inhabitant of xeric habitats in the southwestern United States and northwestern Mexico (Tuskes 1984). Larvae emerge from diapausing egg masses in the spring and feed on Flat-top Buckwheat, *Eriogonum fasciculatum* Bentham (Polygonaceae) (Stone & Smith 1990). The species is univoltine, flying in autumn, with peak emergence from September through early November (Tuskes & McElfresh 1995). Most adults emerge the same year they pupate; however, reared individuals from San Diego County, California have emerged four and one half years after pupation (pers. obs., Powell 1987). Adults are diurnal, non-feeding, and brightly colored (Ferguson 1971, Tuskes et al. 1996). Mate location is facilitated by an airborne pheromone from 'calling' females, and once a female has mated, she stops releasing the pheromone (Tuskes et al. 1996). While investigating aspects of patch-size distribution of this species in southern California, I observed previously unreported mating behavior, on which I report here.

Field work was conducted between 1030 and 1530 PST, 6–19 October, 1996, at Naval Air Station Miramar (parcel G) in San Diego County, California. 11 mm long rubber lures infused with a chemical blend that replicates the primary components of *Hemileuca electra* female pheromone (Jocelyn Millar & Steve McElfresh, unpubl. data) were deployed to attract conspecific males. The dull red-brown lures were kept in a cooler until trials began. In order to observe male response only to their physical presence, two non-calling, sedentary females that had mated on an earlier day were placed approximately 12 cm from the lures on a flat surface. Fifteen males were allowed to land unmolested on the flat sur-




FIG. 1. Male *Hemileuca electra* attempting copulation with a pheromone-releasing rubber lure within 12 cm of two non-calling female *H. electra*.

face where the lure and two non-calling, mated females were resting, and male behavior was recorded. Observations confirm that males are not attracted to the lures if synthetic pheromone has not been added, and females used had attracted males, prior to mating.

Males were attracted within 1–2 min of deploying lures. Fourteen of the 15 males allowed to land did so within 5 cm of the lures, crawled the remaining distance, and attempted copulation with the lure (Fig. 1). These males apparently were not distracted by the presence of non-calling females less than 12 cm away. One male made incidental contact with one of the females as he approached the lure. After crawling around her, he initiated copulation, which the female accepted. I separated the pair approximately 15 seconds later and released the male. He immediately returned, landed near the lure, and attempted copulation with it, ignoring the female. After a few minutes of attempted copulation with the lure, males departed, as did the male that initially contacted the female.

These observations suggest that visual cues do not play a large role in mate location and initiation of copulation in this species. Males are attracted to the source of the pheromone and attempt to mate on contact. Because males ignored females that were 12 cm away, except when incidental contact resulted in female location, color, pattern, or other visual cues in *H. electra* may not play a large role in facilitating mate location.

During the six days of field work, 497 males of *H. electra* were attracted to lures and captured. Three of these lacked a portion of the abdomen, two of which lacked it entirely. These three males were placed in glassine envelopes where they remained alive for approximately 16 h after capture. It is surprising that these males were able to fly, and lived so long following complete or partial abdominal amputation.

When disturbed, adult *Hemileuca* assume a characteristic defensive posture in which the brightly colored abdomen is curled under the thorax to the head (Tuskes et al. 1996). This posture exposes the abdomen to a predator's attack. However, Steve McElfresh (pers. comm.) has observed the Greater Roadrunner, *Geococcyx californianus* Lesson (Neomorphidae), consuming large numbers of *Hemileuca* males, apparently undaunted by this display.

This study was conducted under U. S. Navy contract N68711-96-LT-COO48 to the San Diego Natural History Museum. I thank Norris Bloomfield for assistance with field work; Steve McElfresh and Jocelyn Millar for generously sharing synthetic pheromone; and Larry Gall, Felix Sperling, Jerry Powell, two anonymous referees and especially John Brown for helpful comments that greatly improved the manuscript. This research was supported in part by the Margaret C. Walker Fund for teaching and research in systematic entomology, and a Hatch grant to Felix Sperling.

LITERATURE CITED

- FERGUSON, D. C. 1971. Bombycoidea, Saturniidae (part 1). Moths of America north of Mexico, Fascicle 20.2A. E. W. Classey Ltd. and R. B. D. Publ., Inc. 153 pp.
- POWELL, J. A. 1987. Records of prolonged diapause in Lepidoptera. J. Res. Lepid. 25:83-109.
- TUSKES, P. M. 1984. The biology and distribution of California Hemileucinae (Saturniidae). J. Lepid. Soc. 38:281-309.
- TUSKES, P. M. & S. MCELFRISH. 1995. The biology and distribution of *Hemileuca electra* (Saturniidae) populations in the United States and Mexico, with descriptions of two new subspecies. J. Lepid. Soc. 49:49-71.
- TUSKES, P. M., J. P. TUTTLE & M. M. COLLINS. 1996. The wild silkmths of North America. Cornell Univ. Press, Ithaca, New York. 264 pp.
- STONE, S. E. & M. J. SMITH. 1990. Buckmoths (Lepidoptera: Saturniidae: *Hemileuca*) in relation to southwestern vegetation and foodplants. Desert Plants 10, 13-18:23-30.

DANIEL RUBINOFF, *Division of Insect Biology, Department of Environmental Science, Policy, and Management, University of California, Berkeley, California 94720, USA.*

Received for publication 18 August 1997; revised and accepted 6 January 1998.

Journal of the Lepidopterists' Society
52(2), 1998, 214-216

NEW DISTRIBUTIONAL AND FOODPLANT RECORDS FOR TWENTY CUBAN MOTHS

Additional key words: distribution, larval sampling, light traps, *Estigmene acrea*.

The most recent treatment of the Cuban insect fauna is that of Bruner et al. (1975), which discussed a number of lepidopteran species of economic interest. The purpose of the present paper is to expand upon this base of knowledge, and provide new distributional and foodplant records for 20 species of Lepidoptera from Cuba. All records discussed in the text and Table 1 derive from field collections made since 1990. Larvae were removed from their wild hosts and reared individually to adults in the laboratory in petri dishes, with fresh pieces of foodplant provided daily. Voucher specimens of adults are deposited in the Entomological Collection of the Centro de Investigaciones de Medio Ambiente (CIMA) in Camagüey, Cuba.

Table 1 summarizes the rearing results for 15 moth species. In addition, foodplant relationships were determined for 4 species for which no previous Cuban data were available. These 4 species are discussed in greater detail below, as is the recent capture of *Estigmene acrea*, apparently a new record for the island of Cuba.